

Measuring Weight-for-Length

Measuring Weight

Infants should be weighed using a hospital-grade platform scale. This may be a beam balance scale or a digital (electronic load cell or strain gauge) scale. Check your equipment regularly to make sure you are getting accurate measurements. Scales should be calibrated on a routine basis. Calibration involves putting known weight on the scale to check accuracy. Be sure the scale is placed on a flat, uncarpeted floor.

Procedure

1. Remove shoes, clothing, and diaper from the infant.
2. Place the scale in the "zero" position before you place the infant on the scale.
3. Make sure the child is on the center of the platform.
4. Record the measurement to the nearest decimal fraction.
5. Remove the child from the scale.

Recheck any measurement that is a large change – up or down from previous measurements.



Measuring Length^{1,2}

Best Practice: A platform with an attached yardstick, a fixed head plate, and a movable footplate is required. The footplate can be adjusted so it comes up to the bottom of the infant's heels. This apparatus should be used on a flat surface and requires two people to operate.

Procedure

1. Remove shoes, clothing, and diaper from the infant.
2. Lay the child on the platform.
3. Have one person hold the head of the infant.
4. The other person should keep the infant's knees straight and bring the adjustable footplate up to the infant's heels.
5. Secure the footplate.
6. Remove the infant from the surface.
7. Record the measurement on the yardstick to the nearest 0.1 cm.

Common Practice: Many clinicians measure infants by laying the patient on the paper covering the exam table and marking the positions of the head and the feet on the paper. They then remove the patient and use a measuring tape to quantify the distance between the two pen markings. While this procedure can be very inaccurate due to the incorrect positioning of the infant, movement and crumpling of the paper, and failure to get perpendicular markings by the pen, there are a few tips to getting good length data if this method is used in your office.

Procedure

1. Ask the caregiver who is with the patient to hold the patient as still as possible.
2. Measure the length three times and use the average.
3. If you notice a leveling off or a decline in the patient's length consider a more precise measurement such as the best practice noted above.

References:

1. Lifshitz, Fima. Pediatric Endocrinology Fifth Edition: Volume 2. Growth, Adrenal, Sexual, Thyroid, Calcium and Fluid Balance Disorders. 2007: 4-6.
2. Wales, JKH, Rogol AD, Maarten W., Color J. Atlas of Pediatric Endocrinology and Growth. 1996: 2-3.