

Smart phrases for ED Buprenorphine

.EDBUPMDM

This is a *** year old who presents with concerns of opioid withdrawal in the setting of likely chronic opioid use and dependence. COWS is *** which {DOES / DOES NOT:28277} meet criteria for treatment.

Urine is (positive/negative) for methadone.

Patient {IS/IS NOT:24494} pregnant.

There {IS/IS NO:20953} evidence of significant alcohol or benzodiazepine use disorder.

The patient {DOES / DOES NOT:28277} have any medical or psychiatric contraindications to initiation of buprenorphine in the emergency department.

The patient received (buprenorphine/naloxone/buprenorphine) for a total of ***mg. Final COWS of ***, we {were/not:15343} able to get control of their withdrawal.

They tolerated treatment well with no obvious adverse events.

The patient will follow-up as set up by care management. Patient {will/will not:27381} receive a bridge prescription. They are given and express understanding of specific return precautions and information.

.EDVIROLAB (This is to be used if virology labs are being ordered in the ED that will not result while patient is in the ED. Please let patient know that labs will be referred to VTC and they will follow up with the patient. Please be sure we have updated contact information)

The patient is told that the virology labs that were drawn today will be reviewed by the Virology Treatment Center/MMP Gilman Clinic. They will be contacted by the VTC/MMPGC with follow-up and treatment as indicated. The patient expresses understanding and is in agreement with the referral and plan.